



Research Models and Services

Cotton Rats

Cotton rat

Origin

Their habitat extends from Central America to Southern North America.

Hsd:Cotton Rat

Developed by the National Institutes of Health, Bethesda, Maryland, and Virion Systems, Inc. In 1996, Harlan Laboratories obtained a breeding nucleus from Virion Systems, Inc. Harlan became Envigo in 2015.

Characteristics

The Cotton Rat (*Sigmodon spp.*) belongs to the family Cricetidae. For over half a century the Cotton Rat has served and is still being used as a model for the study of an extensive list of human and rodent pathogens. Currently, its use is most important in advance studies of:

- Human respiratory syncytial virus (RSV) – the primary cause of infant pneumonia (Prince *et al.*, 1979);
- Parainfluenza (PI3) (Murphy *et al.*, 1981; Porter *et al.*, 1991);
- Adenovirus (Pacini DL *et al.*, 1984);
- Other experiments: Pineal gland study; Environmental toxicology; Human genetic therapy using Adenovirus vector.

Reproduction

Females are monogamous. Animals for breeding should be paired at time of weaning (three-four weeks), as pairing at older ages results in increased fighting. The gestation period is about 27 days with a litter size of about five-six. Cotton rats are good foster mothers: they will readily accept foster infants, even if of a different age than their own litter.

Handling

Cotton Rats are not aggressive. However, they will attempt to bite when they are picked up. Further more, since they move very fast and can jump vertically over 30 cm, they are difficult to handle.

Using a tin as environmental enrichment, the animal is hiding in the tin and one can move the tin with the cotton rat to a clean cage.

References

1. Burgdorfer W, *et al.* (1987) Susceptibility of the hispid Cotton Rat (*Sigmodon hispidus*) to Lyme disease spirochete (*Borrelia burgdorferi*). *Am. J. Trop. Hyg.* 37, 624-628.
2. Donnelly TM and Quimby FW (2002) Biology and diseases of other rodents. In: *Laboratory Animal Medicine 2nd edition* (Fox JG, Cohen BJ, Lew FM and Quimby FW, eds), Academic Press, pp 247-307.
3. Faith RE, *et al.* (1997) The Cotton Rat in Biomedical Research. *Lab. Anim. Sci.* 47, 337-345.
4. Ohwada K, Ito T and Katahira K (1994) Reference values for blood chemistry in the cotton rat (*Sigmodon hispidus*). *Scand. J. Lab. Anim. Sci.* 21, 29-31.
5. Ohwada K and Katahira K (1993) Indirect measurement for body surface area of cotton rats. *Exp. Anim.* 42, 635-637.
6. Pacini DL, *et al.* (1984) A new animal model for human respiratory tract disease due to adenovirus. *J. Infect. Dis.* 150, 92-97.
7. Porter DD, *et al.* (1991) Pathogenesis of Human Parainfluenza Virus 3 infection in two species of Cotton Rats: *Sigmodon hispidus* develops bronchiolitis, while *Sigmodon fulviventer* develops interstitial pneumonia. *J. Virol.* 65, 103-111.
8. Prince GA, *et al.* (1978) The pathogenesis of Respiratory Syncytial Virus infection in Cotton Rats. *Am. J. Pathol.* 93.
9. Prince GA, Potash L, Horswood RL, *et al.* (1979) Intramuscular inoculation of live Respiratory Syncytial Virus induces immunity in cotton rats. *Infect. Immun.* 23, 723-728.
10. Prince GA *et al.* (1993) Pathogenesis of Adenovirus type 5 pneumonia in Cotton Rats (*Sigmodon hispidus*). *J. Virol.* 67, 101-111.
11. Randolph JC *et al.* (1995) Nutritional requirements for reproduction in the hispid Cotton Rat (*Sigmodon hispidus*). *J. Mammol.* 76, 1113-1126.
12. Robel GL *et al.* (1996) Environmental, age, and sex effects on the Cotton Rat (*Sigmodon hispidus*) hematology. *J. Wildl. Dis.* 32, 390-394.
13. Walker EP, Warnick F, Hamlet SE *et al.* (1975) Cotton rat. In: *Mammals of the world*. 3rd edition, Vol. II. Baltimore: John Hopkins Univ. Press.
14. Ward LE (2001) Handling the Cotton Rat for research. *Lab. Animal.* 30, 45-50.

Contact us

North America 800.793.7287 EU and Asia envigo.com/contactus info@envigo.com

